

PERSISTENT 22-YEAR CYCLE IN SUNSPOT ACTIVITY: EVIDENCE FOR A RELIC SOLAR MAGNETIC FIELD

K. Mursula¹, I. G. Usoskin^{1*}, and G. A. Kovaltsov²

¹ Dept. of Physical Sciences, P.O.Box 3000, FIN-90014 University of Oulu, Finland
phone/fax: +358-8-5531378/5531287, email: *kalevi.mursula@oulu.fi*, *ilya.usoskin@oulu.fi*

² Ioffe Physical-Technical Institute, Politeknicheskaya 26, 194021 St.Petersburg, Russia
phone/fax: +7-812-2479167/2471017, email: *gena.kovaltsov@pop.ioffe.rssi.ru*

Abstract

We use the recently presented group sunspot number series to show that a persistent 22-year cyclicity exists in sunspot activity throughout the entire period of about 400 years of direct sunspot observations. The amplitude of this cyclicity is about 10% of the present sunspot activity level. A 22-year cyclicity in sunspot activity is naturally produced by the 22-year magnetic polarity cycle in the presence of a relic dipole magnetic field. Accordingly, a persistent 22-year cyclicity in sunspot activity gives strong evidence for the existence of such a relic magnetic field in the Sun. The stable phase and the roughly constant amplitude of this cyclicity during times of very different sunspot activity level strongly support this interpretation.

1 INTRODUCTION

Magnetic activity in the Sun is determined by the dynamo mechanism [Babcock, 1961]. However, a weak relic magnetic field may exist in the Sun's interior since its formation [Cowling, 1945]. Sonett [1982, 1983] tried to find a signature of such a relic field in sunspot activity. Assigning a negative sign to odd solar cycles, he fitted the Wolf sunspot series with a model consisting of two harmonics with periods of 22 and 90 years, and found a small negative offset in the running mean of the model during the last 150 years. He suggested that such an offset gives evidence for a relic solar field. However, his results for the earlier period from the 18th century to mid-19th century were not conclusive since the offset changed sign in late 1700's. Because of this reason, Sonett's results, and the implied evidence for a relic magnetic field remained rather unconvincing. More recently, Bravo & Stewart [1995] studied the difference in the Sun's polar coronal field during subsequent minima, claiming evidence for an inclined dipole

relic field. However, the available data covered only two solar cycle minima which is insufficient to allow a statistically significant conclusion [Boruta, 1996].

It has been argued [Levy & Boyer, 1982, Boyer & Levy, 1984] that the solar dynamo with its 22-year magnetic polarity (Hale) cycle must result, in the presence of a dipole relic field, in a 22-year cyclicity of sunspot activity. Accordingly, a 22-year cyclicity in sunspot activity would provide compelling experimental evidence for the existence of a relic field in the Sun. However, no convincing evidence for such a cyclicity has been found in sunspot activity when using the famous Wolf sunspot series. This is probably because of the rather poor quality of Wolf sunspot data before mid-19th century [Hoyt & Schatten, 1998, Wilson, 1998, Letfus, 1999], and because of the large long-term variations of sunspot activity level (the secular Gleissberg cycle). The poor quality of Wolf sunspot data is also a likely reason for the inconclusive results by Sonett [1983] for the early times.

Here we perform a detailed analysis of the 22-year cyclicity using the new group sunspot numbers (GSN). The GSN series, depicted in Fig. 1a, includes all known archival records of sunspots starting from the observations of G. Galilei in 1610, and gives a more correct measure of early sunspot activity than the Wolf sunspot numbers [[Hoyt & Schatten, 1998]. Contrary to Sonett [1982, 1983], we do not fit the data to any model but rather analyze the raw sunspot data directly.

2 22-YEAR CYCLICITY IN SUNSPOT ACTIVITY

An empirical Gnevyshev-Ohl (G-O) rule [Gnevyshev & Ohl, 1948] orders the sunspot cycles to even-odd pairs so that the intensity (sum of sunspot numbers over a cycle) of the odd cycle of a pair exceeds that of the preceding even cycle. However,

*on leave from Ioffe Phys.-Tech.Inst., St.Petersburg, Russia




Figure 1: (a) Monthly group sunspot number series R_g (thin line) with interpolated gaps (sections of thick line). Sunspot cycle numbers are shown in the lower panel together with the location of the Maunder (MM) and Dalton (DM) minima. (b) Sunspot cycle intensities I_{GO} (see Eq. (1)) for odd (grey squares) and even (open circles) sunspot cycles. (c) Intensity differences D between the two sunspot cycles forming a G-O pair. (d) Running standardized and 11-year averaged GSN series I_R (see Eq. (3)). The times of the 22-year cycle maxima before and during the Maunder minimum [Usoskin et al., 2000] are noted by black circles.

the G-O rule in the Wolf sunspot series is only valid since solar cycle 10 and fails for cycle pairs 4-5 and 8-9 [Gnevyshev & Ohl, 1948, Vitinsky et al., 1986, Wilson, 1988, Storini & Sykora, 1997]. In analogy with [Gnevyshev & Ohl, 1948], we define the intensity $I_{GO}(k)$ of the sunspot cycle number k as the normalized sum of sunspot numbers over the cycle:

$$I_{GO}(k) = \frac{1}{132} \sum_{j=J(k)}^{J(k+1)-1} R_g(j), \quad (1)$$

where $R_g(j)$ is the GSN value for the month number j and $J(k)$ is the month starting cycle k . (As cycle minima we use here the minima of the running 12-month average [Mursula & Ulich, 1998].) The 11-year cycle length (132 months) is used as a normalization factor. The monthly GSN series has some gaps until the end of 18th century. Most data gaps are rather short, but a few gaps are longer than one year. The longest data gap of 27 months was during cycle 0. The data gaps were interpolated using a fit to a binomial curve with a window length of 41 months. The interpolation method was tested with an artificial series (noised 11-year sinusoid with gaps), yielding an accuracy better than 5% even for the longest gaps. The $I_{GO}(k)$ series, depicted in Fig. 1b, allows to examine the validity of the G-O rule in the GSN series. Starting from cycle 6, each even cycle (including cycle 8) is followed by a more intense odd cycle, verifying the G-O rule in the GSN series since the Dalton minimum in 1790-1830. Before the Dalton minimum, the G-O rule is also valid in a phase-reversed form whereby an odd cycle is followed by a more intense even cycle. Note that this phase-reversed G-O rule applies even for the time before the Maunder minimum which occurred from 1645 to 1715. (Noting that the time difference between the last maximum in 1639-1640 before the deep minimum, and the first maximum in 1705 after it, was roughly six 11-year cycles, we have numbered the three cycles before the minimum as -12, -11 and -10). As a quantitative measure of the G-O rule we have depicted in Fig. 1c the difference $D(k) = I_{GO}(k+1) - I_{GO}(k)$ between the intensities of the two cycles forming a pair. These differences are roughly constant, about 10-20, throughout most of the GSN interval, including $D(-11)$, the only value before the Maunder minimum. It is important to note that the $D(k)$ -values do not correlate with sunspot activity. The correlation coefficient between $D(k)$ -values and average pair intensities $((I_{GO}(k) + I_{GO}(k+1))/2)$ is -0.09 ± 0.25 for solar cycles -4 to 21.

Note that in 1830-1930 and 1950-1990, when cycle intensities remained roughly constant (see Fig. 1b), the odd cycles were more intense than either of the two neighboring even cycles. Similarly, in 1740-1790 the even cycles were larger than the two neighboring




Figure 2: *FFT power spectra of the $I_R(i)$ -series before the Maunder minimum (1720-1800) and after it (1830-1996).*

odd cycles. This behaviour suggests that a 22-year periodicity in cycle intensity is the underlying pattern behind the G-O rule. However, the visibility of the 22-year cyclicity is hidden at times when the sunspot activity level is rapidly changing, e.g., during the recovery of activity after the Maunder minimum, around the Dalton minimum and at the start of the recent high cycles in 1940's. During these times the G-O rule is valid only in the fixed order of cycle pairs. Therefore, in order to examine the existence of the 22-year periodicity in cycle intensity in more detail we standardized [Jenkins & Watts, 1969] the monthly GSN series by subtracting the long-term (30-year) trend:

$$U_g(i) = R_g(i) - \frac{1}{361} \sum_{j=-180}^{180} R_g(i+j) \quad (2)$$

and suppressed the dominant 11-year Schwabe cycle by calculating the running 11-year average intensity $I_R(i)$:

$$I_R(i) = \frac{1}{132} \left(\sum_{j=-65}^{65} U_g(i+j) + \frac{1}{2} U_g(i \pm 66) \right) \quad (3)$$

The $I_R(i)$ -series (see Fig. 1d) shows a persistent 22-year cyclicity both between the Maunder and Dalton minima and since the Dalton minimum. The FFT power spectra of these two parts of the $I_R(i)$ -series are shown in Fig. 2 separately. The confidence level [Jenkins & Watts, 1969] of the 22-year peak in

the power spectrum of $I_R(i)$ -series is 0.99 for 1720-1800 and better than 0.9999 for 1830-1996. (Note that some power from the Schwabe cycle still remains in the $I_R(i)$ -series forming the second peak with a period slightly less than 10 years.) In agreement with results obtained above for the discrete series (see Fig. 1c), the peak-to-peak amplitude of the 22-year cyclicity in Fig. 1d is roughly constant and about 10-20, corresponding to approximately 10% of the amplitude of recent solar cycles. Also, the correlation coefficient between the monthly $I_R(i)$ -series and the monthly GSN series is roughly zero (0.05 ± 0.04 and 0.07 ± 0.07 for 1830-1996 and 1720-1790, respectively), implying that the 22-year cyclicity is independent of the overall sunspot activity level. We have shown recently that during the Maunder minimum, when the Schwabe cycle was strongly suppressed, the remaining, seemingly sporadic sunspot activity still shows a significant 22-year periodicity [Usoskin et al., 2000]. Accordingly, a persistent 22-year cycle exists in sunspot activity throughout the whole time interval of nearly 400 years of direct solar observations. The sunspot maxima during the Maunder minimum [Usoskin et al., 2000] occur in phase with the maxima of the 22-year cyclicity before and after the minimum (see Fig. 1d). Moreover, the times of maxima of the $I_R(i)$ -series roughly correspond to even cycles before and to odd cycles after the Dalton minimum (see Figs. 1b and 1d), in agreement with the suggested phase reversal in the G-O rule at the Dalton minimum. The 22-year cyclicity was not clearly visible during the Dalton minimum. However, the two maxima in the $I_R(i)$ -series around the Dalton minimum are separated by about 43 years, implying that the 22-year cyclicity has no sizeable phase change around the Dalton minimum. Therefore, the phase reversal in the G-O rule is not related to the 22-year cyclicity but, most likely, to the known phase catastrophe (e.g. [Vitinsky et al., 1986, Kremliovsky, 1994]) of the Schwabe cycle in the beginning of the Dalton minimum.

3 DISCUSSION AND CONCLUSIONS

The idea of a weak dipole relic solar magnetic field gives a natural explanation for the observed 22-year cyclicity. A relic field can, due to the high conductivity in the solar interior, survive in the Sun for a very long time, even over time scales comparable to the solar age [Cowling, 1945]. Due to a strong amplification by the dynamo fluid motions in the convection zone, such a weak constant field can interact with the poloidal/toroidal dynamo field and hence play a considerable role in the formation of a sunspot cycle [Levy & Boyer, 1982, Boyer & Levy, 1984]. The total magnetic field in the convection layer is enhanced when

the dynamo field has a favorable orientation with respect to the relic field, and suppressed during the next sunspot cycle which has an opposite magnetic orientation in accordance with the Hale law. Thus, a constant relic field leads to cyclicity in sunspot activity with the period of the magnetic polarity cycle.

Concluding, we have shown that a persistent 22-year cyclicity indeed exists in sunspot activity throughout the whole time interval of about 400 years of direct solar observations. The 22-year cyclicity in sunspot activity is naturally produced by the 22-year magnetic Hale cycle in the presence of a relic dipole magnetic field. Accordingly, the detected 22-year cyclicity in sunspot activity gives strong evidence for the existence of such a relic magnetic field in the Sun. The stable phase and the roughly constant amplitude of this cyclicity during times of very different sunspot activity level strongly support this interpretation.

Acknowledgements. The financial support by the Academy of Finland is gratefully acknowledged. We appreciate the electronic service of NOAA for an easy access to sunspot data.

References

- Babcock, H. W., 1961, ApJ, 133, 572.
 Boruta N., 1996, ApJ, 458, 832.
 Boyer, D., Levy, E. H., 1984, ApJ, 277, 848.
 Bravo, S., Stewart, G., 1995, ApJ, 446, 431.
 Cliver, E. W., Boriakoff, V., Bounar, K. H., 1999, J. Geophys. Res., 101(A12), 27091.
 Cowling, T. G., 1945, Mon. Not. R. Astron.Soc., 105, 166.
 Gnevyshev, M. N., Ohl, A. I., 1948, Astron.Zh., 25(1), 18 (in Russian).
 Hoyt, D. V., Schatten, K., 1998, Solar Phys., 179, 189.
 Jenkins, G. M., Watts, D. G., *Spectral Analysis and Its Applications*, Holden-Day, San Francisco, 1969.
 Jokipii, J. R., Levy, E. V., 1977, ApJ, 213, L85.
 Kremliovsky, M. N., 1994, Solar Phys., 151, 351.
 Letfus, V., 1999, Solar Phys., 184, 201.
 Levy, E. H., Boyer, D., 1982, ApJ Lett., 254, L19.
 Mursula, K., Ulich, T., 1998, Geophys. Res.Lett., 25, 1837.
 Richardson, I. G., Cane, H. V., Wibberenz, G., 1999, J. Geophys.Res., 104(A6), 12549.
 Sonett, C. P., 1982, Geophys. Res. Lett., 9, 1313.
 Sonett, C. P., 1983, Nature, 306, 670.
 Storini, M., Sykora, J., 1997, Solar Phys., 176, 417.
 Usoskin, I. G., Mursula, K., Kovaltsov, G. A. 2000, A&A, 354, L33.
 Vitinsky, Yu. I., Kopecky, M., Kuklin, G.V. *Statistics of Sunspot Formation Activity* (in Russian), Nauka, Moscow, 1986.
 Wilson, R. M. 1988, Solar Phys., 117, 269.
 Wilson, R. M. 1998, Solar Phys., 182, 217.